

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶ : A01N 43/42, 43/40</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/19824 (43) International Publication Date: 13 April 2000 (13.04.00)</p>
<p>(21) International Application Number: PCT/US99/23274 (22) International Filing Date: 6 October 1999 (06.10.99) (30) Priority Data: 60/103,320 7 October 1998 (07.10.98) US (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): ADAMS, Jerry, L. [US/US]; 611 Forest Road, Wayne, PA 19087 (US). (74) Agents: DINNER, Dara, L. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).</p>		<p>(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report.</p>
<p>(54) Title: NOVEL TREATMENT FOR STROKE MANAGEMENT (57) Abstract Use of heterocyclic substituted imidazole compounds and compositions in the treatment of CNS injuries to the brain.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

NOVEL TREATMENT FOR STROKE MANAGEMENT

FIELD OF THE INVENTION

This invention relates to the novel use of imidazole compounds in the treatment of stroke and stroke management.

10

BACKGROUND OF THE INVENTION

Interleukin-1 (IL-1) and Tumor Necrosis Factor (TNF) are biological substances produced by a variety of cells, such as monocytes or macrophages. IL-1 has been demonstrated to mediate a variety of biological activities thought to be important in immunoregulation and other physiological conditions such as inflammation [See, e.g.,
15 Dinarello et al., Rev. Infect. Disease, 6, 51 (1984)]. The myriad of known biological activities of IL-1 include the activation of T helper cells, induction of fever, stimulation of prostaglandin or collagenase production, neutrophil chemotaxis, induction of acute phase proteins and the suppression of plasma iron levels.

20 There are many disease states in which excessive or unregulated IL-1 production is implicated in exacerbating and/or causing the disease. These include rheumatoid arthritis, osteoarthritis, endotoxemia and/or toxic shock syndrome, other acute or chronic inflammatory disease states such as the inflammatory reaction induced by endotoxin or inflammatory bowel disease; tuberculosis, atherosclerosis, muscle degeneration, cachexia,
25 psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis, and acute synovitis. Recent evidence also links IL-1 activity to diabetes and pancreatic β cells.

Dinarello, J. Clinical Immunology, 5 (5), 287-297 (1985), reviews the biological activities which have been attributed to IL-1. It should be noted that some of these
30 effects have been described by others as indirect effects of IL-1.

Excessive or unregulated TNF production has been implicated in mediating or exacerbating a number of diseases including rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions; sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary
35 sarcoidosis, bone resorption diseases, reperfusion injury, graft vs. host reaction, allograft rejections, fever and myalgias due to infection, such as influenza, cachexia secondary to

infection or malignancy, cachexia, secondary to acquired immune deficiency syndrome (AIDS), AIDS, ARC (AIDS related complex), keloid formation, scar tissue formation, Crohn's disease, ulcerative colitis, or pyresis.

Interleukin-8 (IL-8) is a chemotactic factor first identified and characterized in 1987. IL-8 is produced by several cell types including mononuclear cells, fibroblasts, endothelial cells, and keratinocytes. Its production from endothelial cells is induced by IL-1, TNF, or lipopolysachharide (LPS). Human IL-8 has been shown to act on Mouse, Guinea Pig, Rat, and Rabbit Neutrophils. Many different names have been applied to IL-8, such as neutrophil attractant/activation protein-1 (NAP-1), monocyte derived neutrophil chemotactic factor (MDNCF), neutrophil activating factor (NAF), and T-cell lymphocyte chemotactic factor.

IL-8 stimulates a number of functions in vitro. It has been shown to have chemoattractant properties for neutrophils, T-lymphocytes, and basophils. In addition it induces histamine release from basophils from both normal and atopic individuals as well as lysozomal enzyme release and respiratory burst from neutrophils. IL-8 has also been shown to increase the surface expression of Mac-1 (CD11b/CD18) on neutrophils without de novo protein synthesis, this may contribute to increased adhesion of the neutrophils to vascular endothelial cells. Many diseases are characterized by massive neutrophil infiltration.

IL-1 and TNF affect a wide variety of cells and tissues and these cytokines as well as other leukocyte derived cytokines are important and critical inflammatory mediators of a wide variety of disease states and conditions. The inhibition of these cytokines is of benefit in controlling, reducing and alleviating many of these disease states.

There remains a need for the treatment, and for the prevention of CNS injuries which are related to the ability of compounds which are cytokine suppressive, i.e. compounds which are capable of inhibiting cytokines, such as IL-1, IL-6, IL-8 and TNF.

SUMMARY OF THE INVENTION

This invention relates to the use of substituted imidazole compounds, or pharmaceutical compositions thereof in the treatment of stroke, and stroke management.

The compounds for use herein are described in PCT/US97/09888 filed 6 June 1997, and published 18 December 1997 as WO 97/47618, Liverton et al., now US Patent 5,859,041 granted January 12, 1999, whose disclosures are incorporated herein by reference in their entirety.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is to the novel use of the compounds described in WO 97/47618 for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries.

5 Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 98/47899 (PCT/US98/07831), Dodd et al., for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed
10 susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 98/52937 (PCT/US98/10807), Anantanarayan et al., now US Patent
15 5,932,576, granted August 3, 1999, for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent and patent application.

20 Another aspect of the present invention relates to the novel use of compounds described in WO 99/03837 (PCT/US98/13419), Wachter et al., for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent
25 application.

Another aspect of the present invention relates to the novel use of compounds described in WO 98/47892 (PCT/US98/07910), Beers et al., for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of
30 making pharmaceutical formulations thereof are also contained within the noted patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 99/01449 (PCT/EP98/03930), Revesz et al., for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed
35 susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of

making pharmaceutical formulations thereof are also contained within the noted patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 98/52941 (PCT/US98/11684), Hanson et al., for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 97/05878 (PCT/US96/12922), De Laszlo et al., now US Patent 5,837,719, granted November 17, 1998, for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent and patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 97/16441 (PCT/US96/17324), De Laszlo et al., for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 98/21957 (PCT/US97/21019), Chang et al., for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 97/12876 (PCT/US96/15880), now US Patent 5,717,100, granted February 10, 1998, Selnick et al., for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 97/16426 (PCT/US96/17477), De Laszlo et al., for the treatment, an

acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent application.

5 Another aspect of the present invention relates to the novel use of compounds described in WO 97/05877 (PCT/US96/12917), De Laszlo et al., now US Patent 5,782,778, granted August 11, 1998, for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making
10 pharmaceutical formulations thereof are also contained within the noted patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 97/16442 (PCT/US96/18539), De Laszlo et al., for the treatment, an acute setting, as well as including prophylactic use, in preventing in those individuals
15 deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 98/52940 (PCT/US98/10436), Anantanarayan et al., for the treatment,
20 an acute setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent application.

Another aspect of the present invention relates to the novel use of compounds described in WO 99/20624 (PCT/EP98/06472) Cheng et al., for the treatment, an acute
25 setting, as well as including prophylactic use, in preventing in those individuals deemed susceptible to, various CNS injuries. Synthetic chemistry, dosages, and methods of making pharmaceutical formulations thereof are also contained within the noted patent application.

30 CNS injuries as defined herein include both open or penetrating head trauma, such as by surgery, or a closed head trauma injury, such as by an injury to the head region. Also included within this definition is ischemic stroke, particularly to the brain area.

Ischemic stroke may be defined as a focal neurologic disorder that results from insufficient blood supply to a particular brain area, usually as a consequence of an
35 embolus, thrombi, or local atheromatous closure of the blood vessel. The role of

inflammatory cytokines in this are has been emerging and the present invention provides a mean for the potential treatment of these injuries. Relatively little treatment, for an acute injury such as these has been available.

In addition to the advantage of using a compound which is an inhibitor of cytokine suppressive binding protein activity (CSBP) for the inhibition of IL-1 or TNF alone which is neuroprotective, inhibition of both cytokines provides for increased efficacy. These compounds may also be useful for all types of strokes, regardless of the thrombotic/hemorrhagic variants. This would allow for early intervention, possible without use of CAT scans, and potential use in conjugation with tPA, or streptokinase, for instance.

TNF- α is a cytokine with proinflammatory actions, including endothelial leukocyte adhesion molecule expression. Leukocytes infiltrate into ischemic brain lesions and hence compounds which inhibit or decrease levels of TNF would be useful for treatment of ischemic brain injury. See Liu et al., Stoke, Vol. 25., No. 7, pp. 1481-88 (1994) whose disclosure is incorporated herein by reference.

Models of closed head injuries and treatment with mixed 5-LO/CO agents is discussed in Shohami et al., J. of Vasc & Clinical Physiology and Pharmacology, Vol. 3, No. 2, pp. 99-107 (1992) whose disclosure is incorporated herein by reference. Treatment which reduced edema formation was found to improve functional outcome in those animals treated.

As noted, these compounds are useful as cytokine inhibitors, and in particular the preferred method of inhibition is the inhibition of the CSBP/p38/RK kinase pathway. A description of the assay for inhibition of the cytokine specific binding protein (CSBP) is also found in WO 95/07922, whose disclosure is incorporated by reference in its entirety, as well as in US Paten 5,777,097. A kinase binding assay is also described herein. Updated versions of a CSBP kinase assay may be found in later filed SB patent applications including WO 98/57966 for instance.

The compounds of WO 97/47618 can be used in the manufacture of a medicament for the prophylactic or therapeutic treatment of any disease state in a human, or other mammal, which is exacerbated or caused by a neurotraumatic event, such as closed head injuries.

In addition, the compounds of WO 99/20624, WO 98/52940, WO 97/16442, WO 97/05877, WO 97/16426, WO 97/12876, WO 98/21957, WO 97/16441, WO 97/05878, WO 98/52941, WO 99/01449, WO 98/47892, WO 99/03837, WO 98/52937, and WO

98/47899, can be used in the manufacture of a medicament for the prophylactic or therapeutic treatment of any disease state in a human, or other mammal, which is exacerbated or caused by a neurotraumatic event, such as closed head injuries.

The compounds of these patent applications are capable of inhibiting
5 proinflammatory cytokines, such as IL-1, IL-6, IL-8 and TNF and are therefore of use in therapy. IL-1, IL-6, IL-8 and TNF affect a wide variety of cells and tissues and these cytokines, as well as other leukocyte-derived cytokines, are important and critical inflammatory mediators of a wide variety of disease states and conditions. The inhibition
10 of these pro-inflammatory cytokines is of benefit in controlling, reducing and alleviating many of these disease states.

Accordingly, the present invention provides for a method of treating a neurotraumatic disease, in a mammal in need thereof, which method comprises administering to said mammal an effective amount of a CSAID™, cytokine suppressive
15 inhibitory compound, wherein the compound is an inhibitor of the CSBP/p38/RK kinase.

As used herein, the term "inhibiting the production of IL-1 (IL-6, IL-8 or TNF)" refers to:

- a) a decrease of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels by inhibition of the *in vivo* release of the
20 cytokine by all cells, including but not limited to monocytes or macrophages;
- b) a down regulation, at the genomic level, of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels;
- c) a down regulation, by inhibition of the direct synthesis of the cytokine (IL-1, IL-6, IL-8 or TNF) as a postranslational event; or
25 d) a down regulation, at the translational level, of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels.

As used herein, the term "cytokine interfering" or "cytokine suppressive amount" refers to an effective amount of a compound of Formula (I) which will cause a decrease in
30 the *in vivo* levels of the cytokine to normal or sub-normal levels, when given to a patient for the prophylaxis or treatment of a disease state which is exacerbated by, or caused by, excessive or unregulated cytokine production.

A new member of the MAP kinase family, alternatively termed CSBP, p38, or RK,
35 has been identified, See Lee *et al.*, Nature, Vol. 300 n(72), 739-746 (1994). Activation of

this novel protein kinase via dual phosphorylation has been observed in different cell systems upon stimulation by a wide spectrum of stimuli, such as physicochemical stress and treatment with lipopolysaccharide or proinflammatory cytokines such as interleukin-1 and tumor necrosis factor. The cytokine biosynthesis inhibitors, of the present invention, compounds of Formula (I), have been determined to be potent and selective inhibitors of CSBP/p38/RK kinase activity. These inhibitors are of aid in determining the signaling pathways involvement in inflammatory responses.

The invention will now be described by reference to the following biological examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention:

In vivo TNF assay:

- (1) Griswold *et al.*, Drugs Under Exp. and Clinical Res., XIX (6), 243-248 (1993); or
(2) Boehm, *et al.*, Journal Of Medicinal Chemistry 39, 3929-3937 (1996)
whose disclosures are incorporated by reference herein in their entirety.

LPS-induced TNF α Production in Mice and Rats

In order to evaluate in vivo inhibition of LPS-induced TNF α production in rodents, both mice and rats are injected with LPS.

Mouse Method

Male Balb/c mice from Charles River Laboratories are pretreated (30 minutes) with compound or vehicle. After the 30 min. pretreat time, the mice are given LPS (lipopolysaccharide from *Escherichia coli* Serotype 055-85, Sigma Chemical Co., St Louis, MO) 25 ug/mouse in 25 ul phosphate buffered saline (pH 7.0) intraperitoneally. Two hours later the mice are killed by CO₂ inhalation and blood samples are collected by exsanguination into heparinized blood collection tubes and stored on ice. The blood samples are centrifuged and the plasma collected and stored at -20°C until assayed for TNF α by ELISA.

Rat Method

Male Lewis rats from Charles River Laboratories are pretreated at various times with compound or vehicle. After a determined pretreat time, the rats are given LPS (lipopolysaccharide from *Escherichia coli* Serotype 055-85, Sigma Chemical Co., St Louis, MO) 3.0 mg/kg intraperitoneally. The rats are killed by CO₂ inhalation and heparinized whole blood is collected from each rat by cardiac puncture 90 minutes

after the LPS injection. The blood samples are centrifuged and the plasma collected for analysis by ELISA for TNF α levels.

ELISA Method

- 5 TNF α levels are measured using a sandwich ELISA, as described in Olivera et al., Circ. Shock, 37, 301-306, (1992), whose disclosure is incorporated by reference in its entirety herein, using a hamster monoclonal antimurine TNF α (Genzyme, Boston, MA) as the capture antibody and a polyclonal rabbit antimurine TNF α (Genzyme) as the second antibody. For detection, a peroxidase-conjugated goat antirabbit antibody
10 (Pierce, Rockford, IL) is added, followed by a substrate for peroxidase (1 mg/ml orthophenylenediamine with 1% urea peroxide). TNF α levels in the plasma samples from each animal are calculated from a standard curve generated with recombinant murine TNF α (Genzyme).

15 LPS-Stimulated Cytokine Production in Human Whole Blood

- Assay: Test compound concentrations are prepared at 10 X concentrations and LPS prepared at 1 ug/ml (final conc. of 50 ng/ml LPS) and added in 50 uL volumes to 1.5 mL eppendorf tubes. Heparinized human whole blood is obtained from healthy
20 volunteers and is dispensed into eppendorf tubes containing compounds and LPS in 0.4 mL volumes and the tubes incubated at 37 C. Following a 4 hour incubation, the tubes are centrifuged at 5000 rpm for 5 minutes in a TOMY microfuge, plasma is withdrawn and frozen at -80 C.

- Cytokine measurement: IL-1 and/or TNF were quantified using a standardized ELISA
25 technology. An in-house ELISA kit was used to detect human IL-1 and TNF. Concentrations of IL-1 or TNF were determined from standard curves of the appropriate cytokine and IC50 values for test compound (concentration that inhibited 50% of LPS-stimulated cytokine production) were calculated by linear regression analysis.

30

CSBP/p38 Kinase Assay:

- This assay measures the CSBP/p38-catalyzed transfer of ³²P from [α -³²P]ATP to threonine residue in an epidermal growth factor receptor (EGFR)-derived peptide (T669) with the following sequence: KRELVEPLTPSGEAPNQALLR (residues 661-
35 681). (See Gallagher *et al.*, "Regulation of Stress Induced Cytokine Production by

Pyridinyl Imidazoles: Inhibition of CSBP Kinase", BioOrganic & Medicinal Chemistry, 1997, 5, 49-64).

Reactions are carried in round bottom 96 well plate (from Corning) in a 30 ml volume. Reactions contained (in final concentration): 25 mM Hepes, pH 7.5; 8 mM
5 MgCl₂; 0.17 mM ATP (the K_m[ATP] of p38 (see Lee et al., Nature 300, n72 pg. 639-746 (Dec. 1994)); 2.5 uCi of [γ-32P]ATP; 0.2 mM sodium orthovanadate; 1 mM DTT; 0.1% BSA; 10% glycerol; 0.67 mM T669 peptide; and 2-4 nM of yeast-expressed, activated and purified p38. Reactions are initiated by the addition of [γ-32P]Mg/ATP, and incubated for 25 min. at 37 °C. Inhibitors (dissolved in
10 DMSO) are incubated with the reaction mixture on ice for 30 minutes prior to adding the 32P-ATP. Final DMSO concentration was 0.16%. Reactions are terminated by adding 10 ul of 0.3 M phosphoric acid, and phosphorylated peptide is isolated from the reactions by capturing it on p81 phosphocellulose filters. Filters are washed with 75 mM phosphoric acids, and incorporated 32P was quantified using beta scintillation
15 counter. Under these conditions, the specific activity of p38 is about 400-450 pmol/pmol enzyme, and the activity linear for up to 2 hr of incubation. The kinase activity values are obtained after subtracting values generated in the absence of substrate which are 10-15% of total values.

20 Prostaglandin endoperoxide synthase-2 (PGHS-2) assay:

This assay describes a method for determining the inhibitory effects of compounds of Formula (I) on human PGHS-2 protein expression in LPS stimulated human monocytes. A suitable assay for PGHS-2 protein expression may be found in a
25 number of publications, including US Patent 5,593,992 whose disclosure is incorporated herein by reference.

TNF-α in Traumatic Brain Injury Assay

This assay provides for examination of the expression of tumor necrosis factor mRNA in specific brain regions which follow experimentally induced lateral fluid-
30 percussion traumatic brain injury (TBI) in rats. Since TNF-α is able to induce nerve growth factor (NGF) and stimulate the release of other cytokines from activated astrocytes, this post-traumatic alteration in gene expression of TNF-α plays an important role in both the acute and regenerative response to CNS trauma. A suitable assay may be found in WO 97/35856 whose disclosure is incorporated herein by reference.

35

CNS Injury model for IL-1 β mRNA

This assay characterizes the regional expression of interleukin-1 β (IL-1 β) mRNA in specific brain regions following experimental lateral fluid-percussion traumatic brain injury (TBI) in rats. Results from these assays indicate that following TBI, the temporal expression of IL-1 β mRNA is regionally stimulated in specific brain regions. These regional changes in cytokines, such as IL-1 β play a role in the post-traumatic pathologic or regenerative sequelae of brain injury. A suitable assay may be found in WO 97/35856 whose disclosure is incorporated herein by reference.

- 10 Evaluation of compounds in specific models for rat focal Ischemic Stroke injury, may be determined by techniques well known to the skilled artisan, as described in some of the publications below. Alternatively, using microsurgical electrocoagulation of the middle cerebral artery under stereotaxic control, a suitable candidate compound is administered by either intravenous, or oral administration.
- 15 Pose 24 hours of MCAO, the neurological deficits and stained forebrain sections are evaluated. Suitable endpoints are 1) neurological deficits in the forelimb, and hindlimb; 2) infarct volume of (mm³); and hemispheric infarct (%; normalized to normal contralateral hemisphere).
- 20 Additional publication which are of value in this invention include, Barone et al., Stroke, 28:6, 1233- 1244 (1997); Stroke, 23: 9, 1337-1347 (1997); Stroke, 25:7, 1481-1488 (1994); Stroke, 24:11, 1746-51 (1993); Stroke, 28:1, 155-162 (1997); Stroke 26:9, 1665-1669 (1995); Cerebrovascular & Brain Metabolism Reviews, 6:4, 341-360 (1994); Brain Research Bulletin, 35:4, 387-392 (1994); Brain Research
- 25 Bulletin, 31: 565-572 (1993); J. Cerebral Blood Flow and Metabolism, 16:3, 260-366 (1996); Neuroscience and Biobehavioral Reviews, 20:3, 445-452 (1996); Cerebrovascular Diseases: 19th Princeton Stroke Conference, Ed. Moskowitz et al., Boston: Butterworth-Heinemann Press, Chapter 7, pp. 75-91 (1995); Immunotherapy in Neuroimmunologic Diseases, Editors, Zhang et al., Martin Dunitz, Ltd. 1998 (pp.
- 30 155-174); Molecular and Chemical Neuropathy, 24: 13-30 (1995); Annals of N.Y. Academy of Sciences, Vol. 825 (1997), Neuroprotective Agents, 3rd International Conference, Editors, Slikker et al., p. 179-193.

35 All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual

publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

5 The above description fully discloses the invention including preferred
embodiments thereof. Modifications and improvements of the embodiments
specifically disclosed herein are within the scope of the following claims. Without
further elaboration, it is believed that one skilled in the art can, using the preceding
description, utilize the present invention to its fullest extent. Therefore, the Examples
10 herein are to be construed as merely illustrative and not a limitation of the scope of
the present invention in any way. The embodiments of the invention in which an
exclusive property or privilege is claimed are defined as follows.

What is Claimed is:

1. A method of treating a CNS injury to the brain, in a mammal in need of such treatment, which method comprises administering to said mammal an effective
5 amount of a cytokine suppressive binding protein compound described in WO 97/47618 which inhibits the CSBP/p38/RK kinase pathway.
2. The method according to Claim 1 wherein the CNS injury is ischemic stroke.
- 10 3. The method according to Claim 1 wherein the CNS injury is caused by surgery, or is an open head injury.
4. The method according to Claim 1 wherein the CNS injury is a closed head injury.
- 15 5. A method of treating a CNS injury to the brain, in a mammal in need of such treatment, which method comprises administering to said mammal an effective amount of a cytokine suppressive binding protein compound described in WO 99/20624, WO 98/52940, WO 97/16442, WO 97/05877, WO 97/16426, WO 97/12876, WO 98/21957, WO 97/16441, WO 97/05878, WO 98/52941, WO
20 99/01449, WO 98/47892, WO 99/03837, WO 98/52937, or WO 98/47899 which inhibits the CSBP/p38/RK kinase pathway.
6. The method according to Claim 5 wherein the CNS injury is ischemic stroke.
- 25 7. The method according to Claim 5 wherein the CNS injury is caused by surgery, or is an open head injury.
8. The method according to Claim 5 wherein the CNS injury is a closed head injury.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/23274

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A01N 43/42, 43/40,

US CL : 514/279, 318, 340, 326, 327, 330, 333

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/279, 318, 340, 326, 327, 330, 333

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN

search terms: P38.USPT AND P38S

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,776,954 A (DE LASZLO et al.) 07 July 1998, entire document.	1-8
Y	US 5,792,778 A (DE LASZLO et al.) 11 August 1998, entire document	1-8

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A document defining the general state of the art which is not considered to be of particular relevance	*X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B earlier document published on or after the international filing date	*Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G document member of the same patent family
*O document referring to an oral disclosure, use, exhibition or other means	
*P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

29 DECEMBER 1999

Date of mailing of the international search report

21 JAN 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

THEODORE J. CRIARES

Telephone No. (703) 308-1235